

# Exact Results for Three-Body Correlations in a Degenerate One-Dimensional Bose Gas

Vadim V. Cheianov,<sup>1</sup> H. Smith,<sup>2</sup> and M. B. Zvonarev<sup>2</sup>

<sup>1</sup>*NORDITA, Blegdamsvej 17, DK-2100 Copenhagen, Denmark*

<sup>2</sup>*Niels Bohr Institute, Universitetsparken 5, DK-2100 Copenhagen, Denmark*

Motivated by recent experiments we derive an exact expression for the correlation function entering the three-body recombination rate for a one-dimensional gas of interacting bosons. The answer, given in terms of two thermodynamic parameters of the Lieb-Liniger model, is valid for all values of the dimensionless coupling  $\gamma$  and contains the previously known results for the Bogoliubov and Tonks-Girardeau regimes as limiting cases. We also investigate finite-size effects by calculating the correlation function for small systems of 3, 4, 5 and 6 particles.

Quantum fluctuations are well known to have a profound influence on the physics of one-dimensional (1D) systems. Due to quantum fluctuations, a 1D gas of weakly interacting bosons does not exhibit true long-range order and hence does not undergo Bose-Einstein condensation at any temperature. With increasing repulsion between the bosons, the gas exhibits a smooth crossover to the strong coupling regime, where the correlations in the positions of the particles endow it with fermion-like thermodynamic properties. A strong repulsion between the bosons also has a marked effect on the local fluctuation properties, leading, for example, to a fermion-like suppression of the fluctuations of particle density. In this paper we shall focus on a three-body correlation function, which is directly related to the lifetime of a Bose gas unstable with respect to three-body recombination processes. It has been studied in a recent experiment, where measurements of the three-body recombination rate for trapped atoms were performed [1].

The idea of using three-body recombination for a measurement of local correlations was originally put forward by Kagan et al. [2]. These authors showed that the low-temperature recombination rate for a 3D Bose gas is proportional to the local three-body correlation function  $g_3$  defined by the ground-state expectation value

$$g_3 = \langle : \hat{n}^3(\mathbf{r}) : \rangle \equiv \langle \hat{\Psi}^\dagger(\mathbf{r})^3 \hat{\Psi}(\mathbf{r})^3 \rangle. \quad (1)$$

Here  $\hat{n}(\mathbf{r}) = \hat{\Psi}^\dagger(\mathbf{r})\hat{\Psi}(\mathbf{r})$  is the particle density operator,  $\hat{\Psi}^\dagger$  ( $\hat{\Psi}$ ) denote boson creation (annihilation) operators and the symbol  $::$  stands for normal ordering. It was noted that, due to the factor-of-six suppression of  $g_3$  in a Bose-Einstein condensate at zero temperature as compared to the non-condensed Bose gas, three-body recombination can be used as a diagnostic tool for distinguishing between the condensed and the non-condensed phases. This was later confirmed experimentally [3].

In the experiment of Ref. [1] the recombination process was used for the investigation of three-body correlations in a 1D Bose gas. A magnetically trapped Bose-Einstein condensate of  $^{87}\text{Rb}$  atoms was loaded into a deep 2D optical lattice, by which the condensate was divided into an array of independent 1D systems. In the case of  $^{87}\text{Rb}$  two-body losses are very small, and it was possible to

model the decay in time of the number of trapped atoms in terms of one-body and three-body processes only. The rate of change of the total number of atoms  $N$  was written as

$$\frac{dN}{dt} = -K_1 N - \int d\mathbf{r} K_3^{1D} n_{3D}^3, \quad (2)$$

where  $K_1$  and  $K_3^{1D}$  are the rate coefficients for one- and three-body losses. By measuring the decrease of the number of atoms as a function of time it was found that the rate coefficient  $K_3^{1D}$  in the 1D gas was reduced considerably, by a factor of about 7, compared to its value  $K_3^{3D}$  for a 3D condensate. The measurements were carried out for a particular value of the dimensionless coupling constant  $\gamma$  ( $\approx 0.5$ ) characterizing the strength of the 1D correlations. Given that  $K_3$  is proportional to the three-body correlation function  $g_3$ , the observed reduction was interpreted as an effect of the reduced dimensionality on  $g_3$ .

In the following we shall therefore focus on calculating  $g_3$  for a 1D Bose gas. Our treatment is based on the Lieb-Liniger (LL) model [4] for which the dimensionless coupling parameter  $\gamma$  is given in Eq. (5) below. In the thermodynamic limit, where the particle number  $N$  and the size of the system  $L$  tend to infinity while the density  $n$  remains constant,

$$n = N/L = \text{const}, \quad N, L \rightarrow \infty, \quad (3)$$

an expression for  $g_3$  has so far been obtained only for small and large values of  $\gamma$  [5]. Here we report an explicit expression for  $g_3$ , valid for all  $\gamma$ . Its derivation employs an integrable lattice regularization of the LL model [6, 7] together with conformal field theory. The calculations are quite involved and will be presented elsewhere [8]. The answer is given by Eq. (17) in terms of two thermodynamic parameters of the LL model: the second- and fourth-order moment, Eq. (16), of the quasi-momentum distribution function. The quasi-momentum distribution function is the solution of the Lieb equation (14), which is a linear integral equation.

Within the LL model the Hamiltonian for  $N$  identical bosons of mass  $m$  is taken to be

$$H = \frac{\hbar^2}{2m} \left[ - \sum_{i=1}^N \frac{\partial^2}{\partial x_i^2} + 2c \sum_{1 \leq i < j \leq N} \delta(x_i - x_j) \right]. \quad (4)$$

The interaction constant  $c \geq 0$  has the dimension of inverse length. The dimensionless coupling strength  $\gamma$  is given by

$$\gamma = c/n. \quad (5)$$

The LL model allows for an exact determination of the eigenfunctions and eigenvalues of  $H$ . In the  $\gamma \rightarrow \infty$  limit of the model, known as the Tonks-Girardeau (TG) gas [9], the eigenvalues are the same as that of a free fermion gas.

While the eigenfunctions and eigenvalues are known exactly for the LL model, its correlation functions in general and  $g_3$  in particular have been much less investigated. The reason for this is that the eigenfunctions are very complicated for a general value of  $N$ . We exhibit these below and calculate  $g_3$  analytically for the simplest possible case  $N = 3$ .

The eigenfunctions  $\Psi(x_1, x_2, \dots, x_N)$  of the Hamiltonian (4) must be symmetric under any interchange of coordinates. Since the particles interact via a delta-function potential, the derivatives of  $\Psi$  jumps when two particles approach each other, while when all  $x_j$  are different,  $\Psi$  satisfies the free-particle Schrödinger equation. The explicit expression for  $\Psi$  was obtained in Ref. [4] and given in Ref. [10] in the following form

$$\Psi(x_1, x_2, \dots, x_N) = C \sum_P (-1)^{[P]} \exp \left\{ i \sum_{j=1}^N k_{P_j} x_j \right\} \times \prod_{j>l} [k_{P_j} - k_{P_l} - ic \operatorname{sgn}(x_j - x_l)], \quad (6)$$

where  $P$  is a permutation of  $N$  numbers and  $[P]$  is the parity of the permutation. The possible values of the quasi-momenta  $k_j$  are determined by the boundary conditions imposed on the system. For periodic boundary conditions (that is, for  $N$  particles placed on a ring of circumference  $L$ )  $k_j$  are solutions to the following system of coupled nonlinear equations, called the Bethe equations,

$$\exp \{ i k_j L \} = \prod_{l \neq j}^N \frac{k_j - k_l + ic}{k_j - k_l - ic}, \quad j = 1, \dots, N. \quad (7)$$

The calculation of the normalization constant  $C$  in Eq. (6) is a nontrivial task. For the periodic boundary conditions a closed expression for  $C$  was suggested by Gaudin (see [11] and references therein) and proved in Ref. [12]; a detailed discussion can be found in Ref. [10]. One has now all the ingredients for calculating  $g_3 \equiv g_3(\gamma, N)$  for the 1D system. Written in the first-quantized form,  $g_3(\gamma, N)$  is

$$g_3(\gamma, N) = \frac{N!}{(N-3)!} \int_0^L dX |\Psi(0, 0, 0, x_4, \dots, x_N)|^2, \quad (8)$$




Figure 1: The three-body correlation function  $g_3$ , normalized to its value in the absence of interactions, as a function of  $\gamma$ . The dashed lines are for particle numbers  $N = 3, 4, 5$ , and  $6$  (from top to bottom), and the full line is for  $N = \infty$ .

where  $dX = dx_4 \dots dx_N$  and  $\Psi$  is given by Eq. (6). Note that the average  $\langle \dots \rangle$  in Eq. (1) is taken over the ground state of the system; equivalently, it is the ground-state wave function which enters Eq. (8). This is ensured by selecting the proper set of the quasi-momenta  $k_j$  among all the solutions of the Bethe equations (7).

All quasi-momenta  $k_j$  are equal to zero for the non-interacting system,  $\gamma = 0$ , and the wave function (6) is uniform in space,  $\Psi = (1/\sqrt{L})^N$ , which implies that

$$g_3(0, N) = n^3 N(N-1)(N-2)/N^3. \quad (9)$$

We now use the wave function (6) to get  $g_3(\gamma, N)$  for  $N = 3$  analytically, and for  $N = 4, 5$  and  $6$  numerically. The quasi-momenta in the ground state of the system with  $N = 3$  obey  $k_1 = -k_3$  and  $k_2 = 0$ . It is convenient to write the Bethe equation for  $k_1$  in the form

$$\lambda = 2\pi - 2 \arctan(\lambda/3\gamma) - 2 \arctan(2\lambda/3\gamma), \quad (10)$$

where we have introduced  $\lambda \equiv k_1 L$  and the principal branch for arctangent has been chosen:  $-\pi/2 \leq \arctan(x) \leq \pi/2$ . The solution to this transcendental equation grows monotonically from  $\lambda \simeq 3\sqrt{\gamma}$  for  $\gamma \ll 1$  to  $\lambda \simeq 2\pi - 4\pi/\gamma$  for  $\gamma \gg 1$ . The resulting answer for  $g_3(\gamma, 3) = 6|\Psi(0, 0, 0)|^2$  is

$$g_3(\gamma, 3) = \frac{16}{3} \frac{A^2 B \lambda^6}{\gamma^3 L^3 [1 + 4(3A^2 + 2A + B + 6AB)]}, \quad (11)$$

where  $A = 3\gamma/(9\gamma^2 + \lambda^2)$  and  $B = 3\gamma/(9\gamma^2 + 4\lambda^2)$ . The ratio  $g_3(\gamma, 3)/g_3(0, 3)$  is plotted in Fig. 1. It decreases monotonously with increasing  $\gamma$ , from  $1 - 29\gamma/24$  for  $\gamma \ll 1$  to  $512\pi^6/243\gamma^6$  for  $\gamma \gg 1$ . In Fig. 1 we also plot  $g_3(\gamma, N)$  calculated for  $N = 4, 5$  and  $6$ . Note that

the magnitude of the slope at the origin ( $\gamma=0$ ) increases with increasing number of particles, and when  $N \rightarrow \infty$  it approaches infinity, in agreement with the Bogoliubov result given in Ref. [5]:

$$g_3(\gamma, \infty)/n^3 \simeq 1 - 6\sqrt{\gamma}/\pi, \quad \gamma \rightarrow 0. \quad (12)$$

We have obtained an analytic expression for  $g_3$  in the Tonks-Girardeau limit,  $\gamma \rightarrow \infty$ , for arbitrary values of  $N$  by generalizing the result Eq. (16) of Ref. [5], valid in the thermodynamic limit, to the case of a finite number of particles. The result is

$$\frac{g_3(\gamma, N)}{n^3} \simeq \frac{16\pi^6}{15\gamma^6} \frac{(N^2 - 1)^2(N^2 - 4)}{N^6}, \quad \gamma \rightarrow \infty. \quad (13)$$

We now present our main result: the exact expression for the three-body correlation function  $g_3(\gamma, \infty)$  in the thermodynamic limit (3) of the Lieb-Liniger model (4). The system of Bethe equations (7) reduces in this limit to the linear integral equation called the Lieb equation:

$$\sigma(k) - \frac{1}{2\pi} \int_{-1}^1 dq \frac{2\alpha\sigma(q)}{\alpha^2 + (k - q)^2} = \frac{1}{2\pi}, \quad (14)$$

where  $\alpha$  is an implicit function of  $\gamma$ :

$$\alpha = \gamma \int_{-1}^1 dk \sigma(k). \quad (15)$$

In terms of the moments  $\epsilon_m$  defined by

$$\epsilon_m = \left(\frac{\gamma}{\alpha}\right)^{m+1} \int_{-1}^1 dk k^m \sigma(k), \quad m = 2, 4. \quad (16)$$

the exact expression for  $g_3(\gamma, \infty)$  is

$$\begin{aligned} \frac{g_3(\gamma, \infty)}{n^3} &= \frac{3}{2\gamma} \epsilon'_4 - \frac{5\epsilon_4}{\gamma^2} \\ &+ \left(1 + \frac{\gamma}{2}\right) \epsilon'_2 - 2\frac{\epsilon_2}{\gamma} - \frac{3\epsilon_2\epsilon'_2}{\gamma} + \frac{9\epsilon_2^2}{\gamma^2}, \end{aligned} \quad (17)$$

where  $\epsilon'_m$  is the derivative of  $\epsilon_m$  with respect to  $\gamma$ .

Equations (14)–(17) form a closed set of equations determining  $g_3(\gamma, \infty)$ . We plot the result in Figs. 1 and 2. For a numerical solution of the integral equation (14) we have used the **Mathematica** package **NISolve** [13]. In Fig. 1 we compare the normalized function  $g_3(\gamma, \infty)/g_3(0, \infty)$  with that calculated for  $N = 3, 4, 5$ , and 6. One can see that the convergence to the thermodynamic limit is quite slow. In Fig. 2 we compare  $g_3(\gamma, \infty)/n^3$  with the experimental data from Ref. [1]. The average value of  $\gamma$ , measured in [1],  $\gamma_m \approx 0.45$ , and the corresponding value of  $g_3(\gamma_m, \infty) \approx 0.14$  are shown as the dot inside the box which represents the experimental uncertainty according to Ref. [1]:  $0.34 < \gamma_m < 0.65$  and  $0.05 < g_3(\gamma_m, \infty) < 0.23$ . The dashed lines show the asymptotic expressions given in Eqs. (12) and (13) for




Figure 2: Double-logarithmic plot of the three-body correlation function  $g_3(\gamma, \infty)$  as a function of  $\gamma$  (full line). The dashed lines indicate the asymptotic results [5] given in Eqs. (12) and (13) for small and large  $\gamma$ , respectively. The measured value is shown as the dot inside a box indicating the experimental uncertainty [1].

small and large  $\gamma$ , respectively. Evidently, the asymptotic expressions do not account for the observed value, and the exact expression (17) is needed.

Eq. (17) gives  $g_3(\gamma, \infty)/n^3 \simeq 16\pi^6/15\gamma^6$  when  $\gamma \rightarrow \infty$ , thus reproducing the asymptotic expression Eq. (13) taken at  $N = \infty$ . To check this is a straightforward task since Eq. (14) admits a regular perturbative expansion with  $1/\alpha$  being a small parameter. The opposite limit,  $\gamma \rightarrow 0$ , of Eq. (17) is much more difficult to analyze since the kernel in Eq. (14) becomes singular when  $\alpha \rightarrow 0$ . Beyond the leading order, the results were obtained only recently [14]:

$$\sigma(k) \simeq \sqrt{1 - k^2}/2\pi\alpha + f(k), \quad \alpha \rightarrow 0 \quad (18)$$

where  $f(k)$  is written explicitly in Ref. [14], see Eq. (4.11). For the moments Eq. (16) one gets  $\epsilon_2 \simeq \gamma(1 - 4\sqrt{\gamma}/3\pi)$  and  $\epsilon_4 \simeq 2\gamma^2(1 - 44\sqrt{\gamma}/15\pi)$ . With these expressions for the moments, expansion (12) is reproduced.

A useful representation of our result (17) is the following approximate form

$$\frac{1 - 6\pi^{-1}\gamma^{\frac{1}{2}} + 1.2656\gamma - 0.2959\gamma^{\frac{3}{2}}}{1 - 0.2262\gamma - 0.1981\gamma^{\frac{3}{2}}}, \quad 0 \leq \gamma \leq 1, \quad (19)$$

$$\frac{0.705 - 0.107\gamma + 5.08 \cdot 10^{-3}\gamma^2}{1 + 3.41\gamma + 0.903\gamma^2 + 0.495\gamma^3}, \quad 1 \leq \gamma \leq 7, \quad (20)$$

$$\frac{16\pi^6}{15\gamma^6} \frac{9.43 - 5.40\gamma + \gamma^2}{89.32 + 10.19\gamma + \gamma^2}, \quad 7 \leq \gamma \leq 30. \quad (21)$$

The relative error of this approximation does not exceed  $2 \cdot 10^{-3}$  for all values of  $\gamma$  in the interval  $0 \leq \gamma \leq 30$ , which we expect to be experimentally relevant. Note that the

asymptotic result (13) for  $N = \infty$  is nearly a factor of two larger than the exact result for  $\gamma = 30$ .

The formal derivation of Eq. (17) is quite lengthy and will be given elsewhere [8]. Here we shall only summarize the main steps: We are interested in the thermodynamic limit of the model, defined by Eq. (3). In this limit it is natural to use a field-theoretical approach rather than the first-quantized formalism we employed for the few-particle systems. The Lieb-Liniger model is completely integrable. In the field-theoretical language complete integrability implies the existence of an infinite family of conserved currents  $\mathcal{J}_n$ ,  $n = 0, 1, 2, \dots$ . The integrals over space of these conserved currents commute with each other, including the Hamiltonian (4). There is thus an infinite set of mutually commuting operators  $J_n$ ,  $n = 0, 1, 2, \dots$ , for which the eigenfunctions and the spectrum are known exactly. A procedure for generating  $J_n$  is discussed, for example, in Ref. [10]. The first three currents generated by this method are  $\mathcal{J}_0 = \mathcal{N}$ ,  $\mathcal{J}_1 = \mathcal{P}$ , and  $\mathcal{J}_2 = \mathcal{H}$ , where  $\mathcal{N}$ ,  $\mathcal{P}$ , and  $\mathcal{H}$  are the number density of particles, the momentum density, and the Hamiltonian density, respectively. These three currents exist for non-integrable models as well, while the currents  $\mathcal{J}_n$  for  $n > 2$  are specific to the LL model. To illustrate our approach we introduce the classical expression for the current  $\mathcal{J}_4$  (its quantization will be discussed in the next paragraph):

$$\mathcal{J}_4 = \Psi^* \partial_x^4 \Psi + c \Psi^* \Psi^* (\partial_x \Psi) \partial_x \Psi + 8c (\partial_x \Psi^*) \Psi^* \Psi \partial_x \Psi + c (\partial_x \Psi^*) (\partial_x \Psi^*) \Psi \Psi + 2c^2 (\Psi^3)^* \Psi^3. \quad (22)$$

The quantum-mechanical version of the last term in Eq. (22) contains  $\hat{\Psi}^\dagger(x)^3 \hat{\Psi}(x)^3$  in which we are interested. When averaged over the ground state, this term gives the desired correlation function  $g_3$ , Eq. (1). Some of the remaining terms in (22) can be eliminated by using the Hellmann-Feynman theorem, which applies to all conserved quantities. This theorem was applied to the Hamiltonian of the LL model in Ref. [5] to calculate the correlation function  $\langle \hat{\Psi}^\dagger(x)^2 \hat{\Psi}(x)^2 \rangle$ . In our case not all terms can be eliminated by use of the Hellmann-Feynman theorem. We have succeeded in obtaining additional identities [8] by employing the long-distance asymptotics of the correlation functions of conserved currents known from the conformal limit of the theory.

The quantization of the classical current Eq. (22) is far from being straightforward. A naive quantum-mechanical version of the expression (22) contains terms involving derivatives of field operators  $\hat{\Psi}$  and  $\hat{\Psi}^\dagger$ . Such terms are not well-defined in general. An explicit demonstration of this was considered in the paper [15], where it was shown that the expectation value  $\langle \hat{\Psi}^\dagger(0) \partial_x^3 \hat{\Psi}(x) \rangle$  is discontinuous at  $x = 0$ . The reason for this behavior can be seen from the structure of the Bethe-ansatz wave function (6), which has a cusp whenever the coordinates of any two particles coincide. This does not

cause difficulties when one considers the momentum operator and the Hamiltonian itself, that is, when one works with the operators  $\hat{\Psi}^\dagger(x) \partial_x \hat{\Psi}(x)$  and  $\hat{\Psi}^\dagger(x) \partial_x^2 \hat{\Psi}(x)$ . However, for other operators we encounter difficulties which we have not been able to resolve within the continuum model (4). Therefore we have used an integrable lattice model, the so-called  $q$ -boson hopping model [6, 7], which reduces to the Lieb-Liniger model (4) in the continuum limit. Within this model all higher (quantum) integrals of motion are defined unambiguously.

In conclusion, we have obtained an exact expression for the three-body correlation function Eq. (1) within the Lieb-Liniger model of a one-dimensional Bose gas. The explicit expression is given by Eq. (17) and Eqs. (19)-(21), and compared in Fig. 2 with a recent experiment. The finite-size effects are explored and a comparison with the thermodynamic limit is given in Fig. 1. We expect that our results will stimulate further experimental investigations of the local correlations in low-dimensional trapped atomic gases.

The authors would like to thank N.M. Bogoliubov for helpful discussions. M.B. Zvonarev's work was supported by the Danish Technical Research Council via the Framework Programme on Superconductivity.

- 
- [1] B. Laburthe Tolra, K.M. O'Hara, J.H. Huckans, W.D. Phillips, S.L. Rolston, and J.V. Porto, Phys. Rev. Lett. **92**, 190401 (2004).
  - [2] Y. Kagan, B.V. Svistunov, and G.V. Shlyapnikov, JETP Lett. **42**, 209 (1985).
  - [3] E.A. Burt, R.W. Ghrist, C.J. Myatt, M.J. Holland, E.A. Cornell, and C.E. Wieman, Phys. Rev. Lett. **79**, 337 (1997).
  - [4] E.H. Lieb and W. Liniger, Phys. Rev. **130**, 1605 (1963).
  - [5] D.M. Gangardt and G.V. Shlyapnikov, Phys. Rev. Lett. **90**, 010401 (2003).
  - [6] N.M. Bogoliubov, R.K. Bullough, and G.D. Pang, Phys. Rev. B **47**, 11495 (1993).
  - [7] N.M. Bogoliubov, A.G. Izergin, and N.A. Kitanine, Nucl. Phys. B **516**, 501 (1998).
  - [8] Vadim V. Cheianov, H. Smith, and M.B. Zvonarev, unpublished.
  - [9] L. Tonks, Phys. Rev. **50**, 955 (1936); M. Girardeau, J. Math. Phys. **1**, 516 (1960).
  - [10] V.E. Korepin, N.M. Bogoliubov, and A.G. Izergin, *Quantum Inverse Scattering Method and Correlation Functions* (Cambridge University Press, Cambridge, England, 1993).
  - [11] M. Gaudin, *La Fonction d'Onde de Bethe*, (Masson, Paris, 1983).
  - [12] V.E. Korepin, Comm. Math. Phys. **86**, 391 (1982).
  - [13] NISolve package is available at <http://library.wolfram.com/infocenter/MathSource/817/>
  - [14] M. Wadati, J. Phys. Soc. Japan **71**, 2657 (2002).
  - [15] Maxim Olshanii and Vanja Dunjko Phys. Rev. Lett. **91**, 090401 (2003).